

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/22, 9/48</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/21439</b> <b>(43) International Publication Date:</b> 18 July 1996 (18.07.96)
<b>(21) International Application Number:</b> PCT/BE96/00002 <b>(22) International Filing Date:</b> 10 January 1996 (10.01.96) <b>(30) Priority Data:</b> 08/370,883 10 January 1995 (10.01.95) US <b>(71) Applicant (for all designated States except US):</b> GALEPHAR P.R. INC. [VC/PR]; Ave Iturregui Calle B, P.O. Box 3468, Carolina, Puerto Rico 00984-3468 (PR). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DEBOECK, Arthur, M. [BE/PR]; HC02 Box 14885, Gurabo, Puerto Rico 00778 (PR). BAUDIER, Philippe [FR/BE]; Avenue Blucher 10, B-1410 Waterloo (BE). MAES, Paul, J. [BE/BE]; Rue Robert Ledecq 8, B-1440 Wauthier-Braine (BE). <b>(74) Agents:</b> SCHMITZ, Y. et al.; Gevers Patents, Holidaystraat 5, B-1831 Diegem (BE).		<b>(81) Designated States:</b> AL, AM, AT, AT (Utility model), AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PHARMACEUTICAL COMPOSITION CONTAINING FENOFIBRATE AND POLYGLYCOLIZED GLYCERIDES		
<b>(57) Abstract</b>  A pharmaceutical composition is provided for treating hyperlipidemia or hypercholesterolemia or both in a mammal, which contains an effective amount of each of fenofibrate and an excipient containing one or more polyglycolized glycerides.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

PHARMACEUTICAL COMPOSITION CONTAINING FENOFIBRATE  
AND POLYGLYCOLIZED GLYCERIDES

BACKGROUND OF THE INVENTION

5 Field of the Invention:

The present invention relates to a pharmaceutical dosage form of fenofibrate having enhanced bioavailability, as well as to an advantageous process for making the same.

Description of the Background:

10 Fenofibrate or p-(4-chlorobenzoyl)-phenoxy isobutyrate isopropyl ester is useful for the treatment of adult patients with very high elevations of serum triglyceride levels and/or cholesterol levels. The usual daily dosage is 300 mg which is administered in two or three doses.

15 Fenofibrate is absorbed as fenofibric acid which is responsible for the pharmacological activity. Fenofibric acid resulting from the hydrolysis of fenofibrate is extensively bound to plasma albumin. The plasma half-life is about 20 hours. Fenofibric acid is excreted

20 predominantly in the urine, mainly as the glucuronide conjugate, but also as a reduced form of fenofibric acid and its glucuronides.

Fenofibrate, is presently available in a pharmaceutical dosage form consisting of hard gelatin

25 capsules containing fenofibrate, lactose starch and magnesium stearate. After oral administration, during a meal, about 60% of the dose of this conventional form is

-2-

effectively absorbed and found in the blood as fenofibric acid, the main metabolite responsible for pharmacological activity. (Strolin & Al, Act Pharmacol. Toxicol. 1986; 59 (Suppl. 5); 167).

5       The first attempt to improve the bioavailability of fenofibrate was performed by Ben-Armor and Al, by solubilizing the fenofibrate in dimethyl isosorbide, a nonaqueous solvent with a miscible wetting agent (Labrafil M 1944CS) with HLB of between 3-4. In order to use the  
10   product in capsules, colloidal silicon oxide was added to increase the viscosity. The liquid so obtained was placed in hard gelatin capsules which, to be leak proof, were sealed. In vivo studies with this formulation indicate that there was no statistically significant difference in  
15   bioavailability between this liquid formulation and the conventional form when the product was given with food.

European Patent Application 0330532 discloses a fenofibrate composition wherein the fenofibrate powder is co-micronized with a solid wetting agent. Sodium lauryl  
20   sulfate is described as the solid wetting agent of choice. The co-micronized powder so obtained is mixed with capsule filling excipient such as lactose, starch, polyvinyl pyrrolidone and magnesium stearate. A formulation of this composition is actually available on the French market  
25   under the trade name Ly pantyl 200 M®. A study comparing this formulation (Ly pantyl 200 M®) to the conventional form

-3-

was undertaken and a statistically significant increase in bioavailability was indicated for the former. In particular, it was found that 67 mg of the new form gives the same amount absorbed as does 100 mg of the conventional form. (J.L. Suichard & Al Cun Therapeutic Research Vol. 54, NS, Nov. 1993).

Unfortunately, co-micronization of the active drug fenofibrate with the wetting agent sodium lauryl sulfate, although necessary, is a time consuming and costly operation. Further, an inherent drawback of micronization is that the material obtained must comply with very stringent particle size specifications.

Moreover, the filling of hard gelatin capsules with a micronized powder is a difficult operation, particularly if weight variation homogeneity is considered.

Hence, a need exists for a fenofibrate formulation that avoids the use of co-micronization, while providing a bioavailability comparable to that afforded by the conventional fenofibrate formulation which uses co-micronization.

#### SUMMARY OF THE INVENTION

Accordingly, it is an object of the present invention to provide a fenofibrate formulation not requiring use of co-micronization which, nevertheless, exhibits a

-4-

bioavailability comparable to formulations of fenofibrate which do.

It is also an object of the present invention to provide a solid, oral dosage form of a fenofibrate formulation that can be prepared by melting the excipients in which the fenofibrate is soluble and, therefore, does not require any particle size specification.

The above objects and others are provided by a pharmaceutical composition for treating hyperlipidemia in and/or hypercholesterolemia a mammal, which contains an effective amount of each of fenofibrate and an excipient containing one or more polyglycolized glycerides.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides a pharmaceutical formulation for treating hyperlipidemia and/or hypercholesterolemia in a mammal, which contains an effective amount of each of a fenofibrate composition and an excipient which contains one or more polyglycolized glycerides, the polyglycolized glycerides preferably having an HLB value of at least about 10.

The present invention is also particularly advantageous for the production of oral solid dosage forms which can be prepared by melting the excipients in which the fenofibrate is soluble, whereby particle size specifications are not required.

-5-

The present invention also relates to the addition of a suspension stabilizer to the molten solution of fenofibrate-polyglycolized glycerides. The suspension stabilizer avoids the formation of fenofibrate crystals during the cooling of the filled hard gelatin capsules. Suitable suspension stabilizers which may be used are, for example, cellulose derivatives, such as hydroxypropylcellulose, hydroxypropylmethyl cellulose, methyl cellulose, and hydroxyethylcellulose, povidone, poloxamers,  $\alpha$ ,  $\Omega$ -hydroxy-poly(oxyethylene) poly(oxypropylene)-poly(oxyethylene) bloc polymers. Other suspension stabilizers equivalent to these stabilizers may, of course, also be used.

The present invention is also particularly advantageous for the production of a pharmaceutical composition in that the hot, homogeneous fenofibrate solution is filled in hard gelatin capsules. This filling process permits the obtention of very precise fenofibrate amounts in each capsule.

The present invention is particularly advantageous as well for the production of the present pharmaceutical composition in that the process for manufacturing the composition requires very few steps such as melting, mixing and filling. This renders the present manufacturing process extremely cost effective when compared to one using co-micronization of powders.

-6-

Polyglycolized glycerides which may be used in the present invention are generally mixtures of known monoesters, diesters and triesters of glycerols and known monoesters and diesters of polyethylene glycols with a mean  
5 relative molecular mass between about 200 and 6000. They may be obtained by partial transesterification of triglycerides with polyethylene glycol or by esterification of glycerol and polyethylene glycol with fatty acids using known reactions. Preferably, the fatty acid component  
10 contains 8-22 carbon atoms, particularly 10-18 carbon atoms. Examples of natural vegetable oils which may be used include palm kernel oil and palm oil. However, these are only examples. The polyol suitably has a molecular weight in the range of about 200-6000 and preferably  
15 contains polyethylene glycol, although other polyols may be employed, such as polyglycerols or sorbitol. They are available on the market under the trade name Gelucire®.

As noted above, the HLB of the polyglycolized glycerides is preferably at least about 10, and more  
20 preferably between about 12 and 15. The melting point of the polyglycolized glycerides may be between about 18°C and 60°C. However, it is especially desirable to use polyglycolized glycerides having a melting point above 30°C, and preferably above 35°C, since there is no need for  
25 sealing the capsule, to assure the leak proofness thereof, when such excipients are used.



-7-

Further, two or more polyglycolized glycerides may be mixed in order to adjust both the HLB value and the melting point to a desired value. The HLB value and melting point of the composition may further be adjusted with the  
5 addition of components such as polyethylene glycols, polyoxyethylene glycols fatty acid esters, and fatty acid alcohols. In view of the present specification, it is well within the skill of the artisan to mix the polyglycolized glycerides to obtain desired HLB values and melting points.

10 It has also been discovered that the present composition affords an increased bioavailability of the fenofibrate as compared to conventional formulations.

Although the present inventors do not wish to be bound by any particular theories, one plausible mechanism of  
15 operation for the present invention is that upon cooling, the melted mixture of hot fenofibrate-polyglycolized glycerides maintains the fenofibrate in liquid form. When absorbed in the gastrointestinal tract of a patient, the gastrointestinal fluids are able to dissolve the  
20 fenofibrate due to the HLB value of the excipient mixture, whereby fenofibrate is readily absorbed.

Generally, the composition of the present invention contains from about 5% to 95% by weight of fenofibrate and from about 95% to 5% by weight of excipient including one  
25 or more polyglycolized glycerides. It is preferred, however, if the present composition contains from about 20%

-8-

to 80% by weight of fenofibrate and from about 80% to 20%  
by weight of excipient. It is even more preferred,  
however, if the present composition contains from about 30%  
to 70% by weight of fenofibrate and from about 70% to 30%  
5 by weight of excipient.

In a particularly preferred composition, generally  
about 45% to 55% by weight of fenofibrate is used and about  
55% to 45% by weight of excipient containing the one or  
more polyglycolized glycerides is used.

10 Generally, the method of the present invention entails  
adding one or more excipients, including the one or more  
polyglycolized glycerides to containing means and then  
heating the excipients until all components are melted.  
Then, fenofibrate is added slowly with continuous stirring  
15 until all fenofibrate added is dissolved. Stirring is then  
continued for about 10 minutes to about 1 hour, and  
preferably for about 15 minutes to about 30 minutes. Then,  
containing means for the pharmaceutical composition, such  
as hard gelatin capsules, are filled with the composition  
20 using a liquid filing capsule machine having dosing pumps  
which are heated to the same temperature as the temperature  
of the molten pharmaceutical composition. Generally, this  
temperature is about 55°C to about 95°C, more typically in  
the range of about 80°C to 90°C. Upon cooling to ambient  
25 temperature, the capsules are packed in bottles. When

-9-

capsules of size 3 are used, each capsule so prepared contains 67 mg of fenofibrate.

It is advantageous, however, to use the following protocol. To about 3 parts by weight polyglycolized glyceride excipient having a melting point of 44°C and an HLB value of 14 molten at 80°C, is added about 2 parts by weight of fenofibrate and about 1 part by weight of hydroxypropyl cellulose. After maintaining the solution under agitation for about 20 additional minutes, hard gelatin capsules are filled therewith.

The present invention will now be further described by reference to certain examples which are provided solely for purposes of illustration and are not intended to be limitative.

15

## EXAMPLE 1

Fenofibrate	6.7 kg
Gelucire® 44/14	5.0 kg
Polyoxamer 407	<u>5.0 kg</u>
	16.7 kg

20

In a stainless steel container, were introduced 5 kg of Gelucire® 44/14 and 5 kg of Poloxamer 407, which were then heated at 85°C until all components are molten. 6.7 kg of fenofibrate was added slowly while continuously stirring the mixture. When all of the fenofibrate was dissolved agitation was maintained for about twenty

25

-10-

minutes. Using a liquid filling capsule machine with dosing pumps heated at 85°C, capsules of size 3 was filled with 167 mg of solution. Upon cooling at room temperature the capsules were packaged in bottles. Each capsule so  
5 prepared contained 67 mg of fenofibrate.

#### PHARMACOKINETICAL STUDY

The composition of Example 1 was compared to conventional form in a pharmacokinetical study with 15 healthy subjects. Each subject received 3 capsules of  
10 composition of Example 1 (201 mg of fenofibrate) or 3 capsules of Lypantyl 100® (300 mg of the conventional form). The sessions were separated by a wash out period of 7 days. The medications were taken after a high-fat  
breakfast. Blood samples were obtained before and at  
15 different times up to 72 hours after administration. The plasma concentration of fenofibric acid was determined in all available samples using a conventional HPLC method.

Plasma Fenofibric Acid Concentration (mg/l vs. time (h) After Administration at 3 Capsules of Example 1 (Total amount of Fenofibrate administered: 201 mg)																	
Post-dose time (h)	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	Mean*	SD
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	0	-
1	BLOQ	BLOQ	0.42	BLOQ	0.52	0.81	0.29	BLOQ	0.12	BLOQ	BLOQ	BLOQ	BLOQ	0.81	BLOQ	0.21	0.30
2	0.36	0.34	3.87	4.31	5.10	6.00	4.66	6.46	2.56	BLOQ	0.99	1.09	3.84	3.03	0.75	2.89	2.19
3	3.31	1.06	7.52	8.12	12.80	7.68	7.50	7.27	6.55	2.51	3.83	3.22	12.68	6.73	5.62	6.43	3.37
4	4.06	2.70	6.02	10.87	13.56	8.27	9.42	8.93	8.16	4.46	5.35	5.23	13.93	7.17	9.61	7.85	3.33
5	4.06	5.49	6.61	10.84	12.65	6.99	9.64	11.70	9.65	6.49	7.42	5.46	14.41	8.53	11.08	8.73	2.99
6	4.32	7.17	6.42	10.68	12.34	6.32	12.19	16.75	11.64	9.75	12.16	5.76	15.68	9.95	13.70	10.32	3.71
7	3.82	7.60	4.28	8.50	11.75	5.68	8.93	8.45	11.43	8.89	11.41	3.74	7.60	9.06	10.72	8.12	2.71
9	4.74	6.83	3.71	6.28	9.61	4.27	8.12	6.19	9.97	6.80	8.79	3.57	7.41	6.42	8.70	6.76	2.05
12	5.61	8.07	2.36	5.66	8.08	3.49	7.05	4.70	7.78	5.00	7.00	6.25	3.75	4.83	6.49	5.74	1.73
24	2.57	3.56	0.85	2.48	4.78	1.39	2.51	1.83	3.48	2.19	2.32	2.30	3.67	2.29	2.64	2.59	0.97
36	1.24	1.53	0.61	1.64	3.01	0.63	1.73	1.16	2.38	1.42	1.64	1.24	1.74	1.26	1.26	1.50	0.61
48	0.80	0.76	0.27	0.98	2.13	0.29	1.05	0.95	1.54	1.06	1.10	0.63	1.33	0.73	0.88	0.97	0.47
60	0.55	0.70	BLOQ	0.64	1.43	0.28	0.73	0.43	0.88	0.73	0.92	0.28	0.78	0.48	0.70	0.64	0.33
72	0.40	0.52	BLOQ	0.50	1.21	BLOQ	BLOQ	0.38	0.68	0.51	0.53	BLOQ	0.62	BLOQ	0.39	0.38	0.34

-12-

Plasma Fenofibric Acid Concentration (mg./l vs. time (h) After Administration at 3 Capsules of the Conventional Form (Total amount of Fenofibrate administered: 300 mg)																
Post- dose time (h)	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	
0	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	
1	BLOQ	BLOQ	BLOQ	0.25	BLOQ	BLOQ	1.90	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	BLOQ	
2	BLOQ	BLOQ	0.25	4.67	0.34	1.52	5.83	BLOQ	BLOQ	0.42	0.63	BLOQ	BLOQ	BLOQ	1.28	
3	1.76	0.99	2.16	7.39	4.51	3.72	5.89	2.45	1.53	1.71	1.55	1.03	1.40	0.47	3.79	
4	3.24	4.62	5.57	9.13	8.83	5.00	5.76	5.12	6.54	4.37	3.58	3.47	4.75	1.48	5.08	
5	4.53	10.24	12.20	12.16	10.43	4.77	6.57	11.97	12.91	4.93	6.94	4.22	6.40	3.55	11.35	
6	8.77	17.36	12.93	12.08	13.18	5.66	6.62	14.17	18.00	9.03	11.45	4.30	11.12	10.65	17.47	
7	4.75	11.92	12.12	10.71	11.36	4.84	5.90	12.31	14.42	8.08	10.58	4.17	13.21	10.11	16.35	
9	3.64	8.21	9.29	8.39	9.62	6.34	5.80	7.33	10.86	6.37	8.25	6.34	10.22	7.21	11.79	
12	4.24	7.03	6.20	6.90	7.96	8.66	5.30	6.67	7.50	5.11	7.09	12.05	9.16	5.74	8.06	
24	2.36	3.43	1.88	3.12	4.76	2.53	2.19	2.61	2.85	2.66	2.85	6.53	4.92	2.29	3.08	
36	1.17	2.03	0.92	1.56	3.27	0.95	1.47	1.14	1.73	1.48	1.38	3.31	2.31	1.33	1.69	
48	0.70	1.17	0.61	1.02	2.06	0.49	0.71	0.94	0.90	1.07	0.92	1.72	1.39	0.81	1.03	
60	0.49	0.50	0.43	0.66	1.77	0.31	0.74	0.81	0.58	0.69	0.55	0.81	1.13	0.54	0.74	
72	BLOQ	BLOQ	0.30	0.49	1.48	BLOQ	0.49	0.54	0.34	0.52	0.40	BLOQ	0.83	0.35	0.40	

-13-

The bioavailability, as measured by the extent of absorption (AUC) indicates, that 3 capsules of Example 1 of the present invention (201 mg of fenofibrate AUC = 195) are bioequivalent to 3 capsules of the conventional form (300 mg of fenofibrate AUC = 221).

That is, the bioavailability of fenofibrate from the composition of Example 1 of the present invention is 1.5 times higher than the bioavailability of fenofibrate of the conventional form.

10

## EXAMPLE 2

Fenofibrate	5 kg
Gelucire® 44/14	7.5 kg
Carbowax 20,000	1.5 kg
Hydroxypropylcellulose	<u>2.5 kg</u>
	16.5 kg

15

To a heated kettle, 7.5 kg of Gelucire® 44/14 and 1.5 kg of carbowax 20,000 were added and then heated at 85°C until all components are molten. 5 kg of fenofibrate was added slowly while continuously stirring. When all the fenofibrate was dissolved, 2.5 kg of hydroxypropylcellulose was added and agitation was maintained for about twenty minutes. Using a liquid filling capsule machine with dosing pumps heated at 85°C, capsules of size 0 were filled with 660 mg of solution. Upon cooling at room temperature the capsules were packaged in bottles. Each capsule so

25

-14-

prepared contained 200 mg of fenofibrate. 12,701 capsules were produced and individually weighed. Results of the capsule weighing is shown in Table 3.

TABLE 3 Capsules Weight Variations From 12,701 Capsules		
5	Theoretical Weight	764.5 mg
	Mean weight of acceptable capsules (95-105%)	763.9 mg
	Standard Deviation of Accepted Capsules	6.9 mg
10	Relative Standard Deviation of Accepted Capsules	0.9%
	Percent of Rejected Capsules (below 95% of Theoretical Amount)	0.307%
15	Percent of Rejected Capsules (above 105% of Theoretical Amount)	0.039%

It may readily be appreciated from Table 3 that the filling process of the present invention is extremely accurate.

#### PHARMACOKINETICAL STUDY

The composition of Example 2 of the present invention was compared during a Pharmacokinetical study to the co-micronized formulation available on the French market (Lypanthyl 200 M®).

The study was conducted as a single dose, randomized, four-way cross over study in 8 healthy subjects. The



-15-

subjects were randomly assigned to one of four administration sequences. On each of the four sessions, separated by wash-out periods of 7 days, the subjects received either 200 mg of fenofibrate under the form  
5 Lypanthyl 200 M® or 200 mg of fenofibrate under the form of Example 2 with and without a high-fat breakfast. Blood samples were taken before and at different times up to 72 hours after administration. The plasma concentrations of fenofibric acid was determined in the samples using on HPLC  
10 Method..

The pharmacokinetics parameters obtained are shown in Table 4.

15

TABLE 4 Pharmacokinetical Parameters After Administration of Lypanthyl 200 M® and Composition of Example 2 Taken With and Without a High Fat Breakfast (Dose 200 mg of Fenofibrate)				
	Without Food		With Food	
	Example 2	Lipanthyl 200M®	Example 2	Lypanthyl 200M®
AUC <sub>0-72</sub>	107.0	101.0	181.0	184.7
C <sub>max</sub>	5.1	5.9	11.1	10.9
T <sub>max</sub>	5.9	5.2	5.2	5.7

20 The present composition may thus be advantageously used to treat hyperlipidemia and/or hypercholesterolemia in humans. Generally, the effective daily amount of fenofibrate from humans ranges from about 100 mg to 600 mg per day, and preferably from about 100 to 300 mg per day,  
25 with the precise amount being determined by the attending

-16-

physician, considering such parameters as condition severity and body weight, for example.

Having fully described the present invention, it will be apparent to one of ordinary skill in the art that many  
5 changes and modification may be made to the above-described embodiments without departing from the spirit and scope of the present invention.

- 17 -

### CLAIMS

1. A pharmaceutical composition for treating hyperlipidemia or hypercholesterolemia or both in a mammal,  
5 which comprises an effective amount of each of fenofibrate and an excipient comprising one or more polyglycolized glycerides.
2. The composition of Claim 1, wherein said fenofibrate is present in an amount of 5% to 95% by weight  
10 based on the total weight of the composition.
3. The composition of Claim 1, wherein the polyglycolized glycerides have a HLB value of at least 10.
4. The composition of Claim 3, wherein the polyglycolized glycerides have a HLB value of from 12 to  
15 15.
5. The composition of Claim 1, which further comprises polyalkylene glycols to adjust the HLB value or melting point or both to the desired value.
6. The composition of Claim 1, wherein a suspension  
20 stabilizer is added.
7. The composition of Claim 6, wherein said suspension stabilizer is selected from the group and consisting of cellulose, povidone, poloxamers,  $\alpha$ ,  $\Omega$ -hydroxy-poly(oxyethylene) poly(oxypropylene)-  
25 poly(oxyethylene) bloc polymers.

-18-

8. The composition of Claim 1, in which said fenofibrate and said excipient are in unit dosage form and are contained in a hard gelatin capsule.

9. The composition of Claim 8, wherein said hard  
5 gelatin capsule contains from about 67 mg to about 200 mg of fenofibrate.

10. A method of making a solid oral dosage form of a pharmaceutical composition, comprising an effective amount of each of fenofibrate and an excipient comprising one or  
10 more polyglycolyzed glycerides, which method comprises adding said molten fenofibrate and said excipient to hard gelatin capsules, and allowing said said molten fenofibrate and said excipient to cool therein.

11. A method of treating hyperlipidemia or  
15 hypercholesterolemia or both in a mammal in need thereof, which comprises administering to said mammal an effective amount of a pharmaceutical composition, comprising fenofibrate and an excipient containing one or more polyglycolyzed glycerides.

20 12. The method of Claim 11, wherein said mammal is human, and said effective amount of fenofibrate in said composition is from about 100 mg to 600 mg per day.

13. The method of Claim 12, wherein said effective  
amount of fenofibrate in said composition is from about 100  
25 mg to 300 mg per day.

-19-

14. The method of Claim 11, wherein said composition is administered orally.

15. The method of Claim 10, which is with the proviso that the fenofibrate used is not co-micronized.

## INTERNATIONAL SEARCH REPORT

 Int. Application No  
 PCT/BE 96/00002

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 A61K31/22 A61K9/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,95 24893 (R. P. SCHERER LTD.) 21 September 1995 see claim 1 see page 13, line 5 - page 15, line 7 see page 25, line 3 - line 4 see page 44; example 6 -----	1-4,6-15

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

29 March 1996

Date of mailing of the international search report

11. 04. 96

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/BE 96/00002

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 11-14 are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

### Information on patent family members

**Internal Application No**

PC1/BE 96/00002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9524893	21-09-95	AU-B- 1897495	03-10-95
-----			





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>A61K 31/22, 9/48</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 96/21439</b> (43) International Publication Date: 18 July 1996 (18.07.96)</p>
<p>(21) International Application Number: <b>PCT/BE96/00002</b> (22) International Filing Date: 10 January 1996 (10.01.96) (30) Priority Data: 08/370,883 10 January 1995 (10.01.95) US (71) Applicant (for all designated States except US): GALEPHAR P.R. INC. [VC/US]; Ave Iturregui Calle B, P.O. Box 3468, Carolina, Puerto Rico 00984-3468 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DEBOECK, Arthur, M. [BE/US]; HC02 Box 14885, Gurabo, Puerto Rico 00778 (US). BAUDIER, Philippe [FR/BE]; Avenue Blucher 10, B- 1410 Waterloo (BE). MAES, Paul, J. [BE/BE]; Rue Robert Ledecq 8, B-1440 Wauthier-Braine (BE). (74) Agents: SCHMITZ, Y. et al.; Gevers Patents, Holidaystraat 5, B-1831 Diegem (BE).</p>		<p>(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i></p>
<p>(54) Title: PHARMACEUTICAL COMPOSITION CONTAINING FENOFIBRATE AND POLYGLYCOLIZED GLYCERIDES (57) Abstract  A pharmaceutical composition is provided for treating hyperlipidemia or hypercholesterolemia or both in a mammal, which contains an effective amount of each of fenofibrate and an excipient containing one or more polyglycolized glycerides.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam